

Report for 2002TX50B: Increase Water Use Efficiency: Implementation of Limited Irrigation for Crop Biotic and Abiotic Stress Management

- unclassified:
 - Falkenberg, Nyland. Site Specific Management of Plant Stress Using Infrared Thermometers and Accu-Pulse. 2002 American Society of Agronomy Meeting (ASA), Indianapolis, IN.
 - Falkenberg, Nyland. Remote Sensing for Site Specific Management of Biotic and Abiotic Stress in Cotton. 2003 Beltwide Cotton Conference, Nashville, TN.
 - Falkenberg, Nyland. Will be presenting again at the Beltwide Cotton Conference in San Antonio, TX and in Denver, CO at the American Society of Agronomy Meeting for the 2003-04 meetings.
 - Falkenberg, N. R., G. Piccinni, M. K. Owens, and J. T. Cothren. Increased Water Use Efficiency-Limited Irrigation to Manage Crop Stress: A Remote Sensing Study. Texas Water Resources Institute SR 2003-003.

Report Follows:

REMOTE SENSING FOR SITE-SPECIFIC MANAGEMENT OF BIOTIC AND ABIOTIC STRESS IN COTTON

Nyland Falkenberg¹, Giovanni Piccinni¹,
M.K. Owens¹, and Dr. Tom Cothren²

¹Texas Agricultural Research and Extension Center
Uvalde, TX

²Texas Agricultural Experiment Station
College Station, TX

Abstract

This study evaluated the applicability of remote sensing instrumentation for site-specific management of abiotic and biotic stress on cotton grown under a center pivot. Three different irrigation regimes (100%, 75%, and 50% ETc) were imposed in the cotton field to: 1) monitor canopy temperatures of cotton with infrared thermometers (IRTs) to pinpoint areas of biotic and abiotic stresses, 2) compare aerial infrared photography to IRTs mounted on center pivots to correlate areas of biotic and abiotic stresses, and 3) to relate yield and yield parameters relative to canopy temperatures. Pivot mounted IRTs and IR cameras were able to differentiate water stress between the irrigation regimes. However, only the IR cameras were effectively able to distinguish between biotic (cotton root rot) and abiotic (drought) stresses with the assistance of ground-truthing. Cooler canopy temperatures were reflected in higher lint yields. The 50% ETc regime had significantly higher canopy temperatures, which were reflected in significantly lower lint yields when compared to the 75 and 100% ETc regimes. Deficit irrigation up to 75% ETc had no impact on yield, indicating that for this year water savings were possible without yield depletion. Canopy temperatures were effective in monitoring plant stress during the canopy development.

Introduction

In 1993, the Texas Legislature placed water restrictions on the farming industry by limiting growers to a maximum use of 2 acre-foot of water per year in the Edwards Aquifer Region. Since then, maximization of agricultural production efficiency has become a high priority for numerous studies in the Winter Garden Area of Texas. Recent investigations have proposed Site-Specific Management (SSM) as an alternative to address this problem. SSM involves satellite-based remote sensing technology and mapping systems to detect specific areas suffering from stress within a field (i.e. water, insect, and disease stress). Crop canopy temperature has been found to be an effective indicator of plant water stress (Moran, 1994). Coupled with remote sensing technology, this concept allows collection and analysis of temperature data from crops using infrared thermometers (IRTs). IRTs mounted on irrigation systems or operated from aircraft can detect water stress by recording changes in leaf temperature caused by the alteration of the soil-plant water flow continuum (Hatfield and Pinter, 1993; Michels et al., 1999). Therefore, remote sensing equipment and mapping systems provide an excellent potential for producers to grow crops under high water use efficiency, by treating only the areas where treatment is needed (i.e. irrigation).

Objectives

The overall objectives of this project are as follow: 1) use remote sensing instrumentation for locating areas showing biotic and abiotic stress signs and/or symptoms in a cotton field, 2) evaluate canopy temperature changes in cotton with the use of IRTs, 3) and assess yield and yield parameters relative to the canopy temperatures.

Materials and Methods

The experiment was conducted at the Texas A&M Agricultural Research and Extension Center in Uvalde, Texas. Cotton variety Stoneville 4892B/Round-up Ready was planted in a circle at 50,000 ppa on 40-inch row spacing and grown under a center-pivot LESA (Low Elevation Sprinkler Application) irrigation system. Furrow dikes were placed between beds to increase water capture and minimize run-off. The soil type is a Knippa clay soil (fine-silty, mixed, hyperthermic Aridic Calciustolls) with a pH of 8.1. Three irrigation regimes (100%ETc, 75%ETc, and 50% ETc) were replicated twice in a randomized block design. A 90-degree wedge was divided equally into six 15-degree regimes, which were maintained at the above mentioned (ETc) values. Thirty Exergen (Irt/c.01-T80F/27C) infrared thermometers (IRTs) were mounted at approximately 15-foot spacing along the pivot length to scan the canopy temperature as the pivot moved. The IRTs recorded canopy temperatures every 10 seconds, and average temperature values every 60 seconds, on a 21X Campbell Scientific datalogger. In addition, canopy temperature differences were determined among treatments using a helicopter equipped with a Mikron 7200 LWIR (Long Wave-length Infrared) infrared camera with infrared band of 8-14 microns. Physiological parameters (i.e., leaf water and osmotic potential) were taken from leaves to determine the level of stress imposed by the different irrigation regimes and the presence of disease. Temperature data were statistically analyzed by ANOVA and separated by Fisher's LSD at $\alpha = 0.05$. Aerial infrared temperature readings were analyzed by using the program Mikroskan 2.6.

Results and Discussion

Environmental conditions for the 2002 cotton season showed that the minimum and maximum temperatures were normal for the area, but excessive rainfall in the month of July prevented the imposition of differential irrigation regimes. Significant differences in canopy temperature were detected in all three irrigation regimes, with a linear increase in canopy temperature resulting from a decrease in plant water availability. Extreme temperatures detected early in crop development were related to the detection of bare soil and moisture availability in the soil by the IRTs. Pivot mounted IRTs were effective in detecting crop canopy temperature differences between the 3 irrigation regimes. Early in the season there were significant differences between all three irrigation regimes; however, at the end of the growing season no significant differences were found between the 100 and 75% ETc regimes. These results are also best explained by yield differences. No significant differences in lint yield were found between the 75% and 100% ETc

regimes. Yield from the 50% regime were significantly less than the 75% and 100% ETc regimes. This yield reduction is associated with increased canopy temperatures of this regime. Yields were 1160 lb/acre, 1420 lb/acre, and 1600 lb/acre for the 50%, 75%, and 100% ETc treatments, respectively. Abiotic and biotic stress can be differentiated better by the Mikron 7200 than the pivot mounted IRTs because of its increased image scanning resolution. The IR camera was able to detect distinct canopy temperature differences between all 3 irrigation regimes. Biotic stress (root rot) was detected by using the camera before symptoms could be detected visually. Pivot mounted IRTs and IR cameras were able to differentiate water stress between the irrigation regimes, but only IR cameras were able to distinguish between abiotic and biotic stress. There was an excellent correlation between canopy temperature and lint yield. Deficit irrigation up to 75% ETc had no impact on yield, indicating that water savings are possible without yield depletion. Also, canopy temperature can be an excellent tool to monitor plant stress.

References

- Barnes, E.M., P.J. Pinter, B.A. Kimball, D.J. Hunsaker, G.W. Wall, and R.L. LaMorte. 2000. Precision irrigation management using modeling and remote sensing approaches. In 4th Decennial Nat. Irri. Symposium. 332-337.
- Bugbee, B., M. Droter, O. Monje, and B. Tanner. 1998. Evaluation and modification of commercial infra-red transducers for leaf temperature measurement. advances in space research. 1-10.
- Burke, J.J., J. L. Hatfield, and D. F. Wanjura. 1990. A thermal stress index for cotton. Agron. J. 82: 526-530.
- Ehler, W.I., S.B. Idso, R.D. Jackson, and R.J. Reginato. 1978. Diurnal changes in plant water potential and canopy temperatures of wheat as affected by drought. Agron. J. 70: 999-1004.
- Grimes, D.W., H. Yamada, and S.W. Hughs. 1987. Climate-normalized cotton leaf water potentials for irrigation scheduling. Agri Water Mgmt. 12: 293-304.
- Hatfield, J.L. 1990. Measuring plant stress with an infrared thermometer. HortScience. 25: 1535-1538.
- Hatfield, J. L., and P.J. Pinter, Jr. 1993. Remote sensing for crop protection. Crop Prot. 12: 403-413.
- Humes, K.S., W.P. Kustas, and M.S. Moran, Use of remote sensing and reference site measurements to estimate instantaneous surface energy balance components over a semiarid rangeland watershed. *Water Resources Research*, Vol. 30, No. 5, pp. 1363-1373, 1994.

- Isdo, S.B., R.J. Reginato, D.C. Reicosky, and J.L. Hatfield. 1982. Determining soil-induced plant water potential depressions in Alfalfa by means of infrared thermometry. *Agronomy Journal*. 73: 826-830.
- Jeger, M. J. and S.D. Lyda. 1986. *Phymatotrichum* root rot (*Phymatotrichum omnivorum*) in cotton: environmental correlates of final incidence and forecasting criteria. *Annals of Applied Biology*. 109: 523-534.
- Jenkins, J.N., J.C. McCarty, and W.L. Parrott. 1990. Competition between adjacent fruiting forms in cotton. *Crop Sci.* 30: 857-860.
- Kenerley, C.M., and M.J. Jeger. 1990. Root colonization by *Phymatotrichum omnivorum* and symptom expression of *Phymatotrichum* root rot in cotton in relation to planting date, soil temperature and soil water potential. *Plant Pathology*. 39: 489-500.
- Kenerley, C.M., T.L. White, M.J. Jeger, and T.J. Gerik. 1998. Sclerotial formation and strand growth of *Phymatotrichopsis omnivora* in minirhizotrons planted with cotton at different soil water potentials. 47: 259-266.
- King, C.J., and H.F. Loomis. 1929. Cotton root rot investigations in Arizona. *Journal Agricultural Research*. 39: 199-221.
- Kustas, W.P., M.S. Moran, K.S. Humes, D.I. Stannard, P.J. Pinter, Jr., L.E. Hipps, E. Swiatek, and D.C. Goodrich, Surface energy balance estimates at local and regional scales using optical remote sensing from an aircraft platform and atmospheric data collected over semiarid rangelands, *Water Resources Research*, Vol. 30, No. 5, pp. 1241-1259, 1994.
- Lyda, S.D., 1978. Ecology of *Phymatotrichum omnivorum*. *Annual Review of Phytopathology*. 16: 193-209.
- Maas, S.J., G.J. Fitzgerald, W.R. DeTar, and P.J. Pinter Jr. 1999. Detection of water stress in cotton using Multispectral remote sensing. *Proc-Beltwide-Cotton- Conf.* 1: 584-585.
- Mahrer, Y. 1990. Irrigation Scheduling with an Evapotranspiration Model: A Verification Study. *Acta Hort.* (ISHS) 278:491-500
- Marani, A., R.B. Hutmacher, and C.J. Phene. 1993. Validation of CALGOS: simulation of leaf water potential in drip-irrigated cotton. *Proc-Beltwide-Cotton-Conf.* 3: 1225-1228.
- Michels, G.J. Jr., G. Piccinni, C.M. Rush, and D.A. Fritts. 1999. Using infrared transducers to sense greenbug (homoptera: aphididae) infestations in winter wheat. *Southwestern Entomologist*. 24: 269-279.

- Moran, M.S., Y. Inoune, and E.M. Barnes. 1997. Opportunities and limitations for image-based remote sensing in precision crop management. *Remote Sens. Environ.* 61: 319-346.
- Nilsson, H.E. 1995. Remote sensing and image analysis in plant pathology. *Annual Review of Phtopathology.* 33: 489-527.
- Plant, R.E., D.S. Munk, B.R. Roberts, R.N. Vargas, R.L. Travis, D.W. Rains, and R.B. Hutmacher. 2001. Application of Remote Sensing to Strategic Questions in Cotton Management and Research. *Journal of Cotton Science.* 5: 30-41.
- Rogers, C.H. 1942. Cotton root rot studies with special reference to sclerotia, covercrops, rotations, tillage, seedling rates, soil fungicides, and effects on seed quality. *Bulletin 614, Texas Agricultural Experiment Station, 45pp.*
- Rush, C.M., D.R. Upchurch, and T.J. Gerik. 1984. In situ observations of *Phymatotrichum omnivorum* with a boroscope minirhizotron system. *Phytopathology.* 74: 104-105.
- Schepers, J.S., and D.D. Francis. 1998. Precision agriculture – what’s in our future. *Commun. Soil Sci. Plant Anal.* 29: 1463-1469.
- Streets, R.B., and H.E. Bloss. 1973. *Phymatotrichum Root Rot.* APS Monograph no. 8., St. Paul, Minnesota, USA: American Phytopathological Society.
- Stewart, R.B., E. I. Mukammal, and J. Wiebe. 1978. The use of thermal imagery in defining frost prone areas in the Niagra fruit belt. *Remote Sens. Environ.* 7: 187-202.
- Wanjura, D.F., D.R. Upchurch, J.R. Mahan. 1992. Automated irrigation based on threshold canopy temperature. *Trans ASAE.* 35: 153-159.
- Yang, C. and G.L. Anderson. 2000. Mapping grain sorghum yield variability using airborne digital videography. *Precision agriculture.* 2: 7-23.

Acknowledgements

Megan Laffere
Cotton Physiology Workgroup at College Station
For additional information email: N-falkenberg@tamu.edu